



Center for
Health Care Strategies, Inc.

FACES OF MEDICAID
DATA SERIES

Bibliography:
**Clarifying Multimorbidity to
Improve Targeting and
Delivery of Clinical Services
for Medicaid Populations –
Literature Review**

December 2010

Cynthia Boyd, MD, MPH*
Bruce Leff, MD*
Carlos Weiss, MD, MHS*
Jennifer Wolff, PhD*
Rebecca Clark
Thomas Richards, MS

The Johns Hopkins University

**Co-Principal Investigators and contributed equally to this report.*

For related materials, visit www.chcs.org.

Made possible through support from Kaiser Permanente.

Overview

This bibliography is part of the *Faces of Medicaid* analysis, *Clarifying Multimorbidity to Improve Targeting and Delivery of Clinical Services for Medicaid Populations*, which was undertaken by the Center for Health Care Strategies (CHCS) and The Johns Hopkins University School of Medicine and Bloomberg School of Public Health. The analysis sought to help policymakers identify intervention strategies with the potential to both improve quality and reduce costs for adult Medicaid beneficiaries with multiple chronic conditions.

In tandem with the analysis, the Johns Hopkins research team performed a literature search to identify, existing clinical opportunities for multimorbid patients. The literature search identified intervention opportunities for general multimorbidity as well as for specific multimorbidity patterns that are outlined in the analysis. Following is a bibliography of the identified literature. For the full analysis and related materials, visit the online resource center at www.chcs.org.

A

- ACCORD group. (2008). The Action to Control Cardiovascular Risk in Diabetes Study Group. *New England Journal of Medicine*, 358, 2545-2559.
- Alexander, J. L., Phillips, W. T., & Wagner, C. L. (2008, May/June). The Effect of Strength Training on Functional Fitness in Older Patients with Chronic Lung Disease Enrolled in Pulmonary Rehabilitation. *Rehabilitation Nursing*, 33(3), 91-98.
- Alexopoulos, G.S., Jeste, D.V., Chung, H., Carpenter, D., Ross, R., & Docherty, J.P. (2005, January). Treatment of dementia and its behavioral disturbances. Introduction: methods, commentary, and summary. *Postgraduate Medicine, Spec No*, 6-22.
- Alexopoulos, G. S., Raue, P. J., Sirey, J. A., & Arean, P. A. (2008, May). Developing an intervention for depressed, chronically medically ill elders: a model from COPD. *International Journal of Geriatric Psychiatry*, 23(5), 447-453.
- Alkema, G.E., Wilber, K.H., Shannon, G.R., & Allen, D. (2007, August). Reduced Mortality: The Unexpected Impact of a Telephone-Based Care Management Intervention for Older Adults in Managed Care. *Health Services Research*, 42(4), 1632-1650.
- Allen, D., Alkema, G. E., Wilber, K. H., & Shannon, G. R. (2007, August). Reduced Mortality: The Unexpected Impact of a Telephone-Based Care Management Intervention for Older Adults in Managed Care. *Health Services Research*, 42(4), 1632-1650.
- Alves, T. C., Rays, J., Telles, R. M., & Fraguas, R. (2007, Jan/Feb). Effects of Antidepressant Treatment on Cognitive Performance in Elderly Subjects With Heart Failure and Comorbid Major Depression: An Exploratory Study. *Psychosomatics*, 48(1), 22-31.
- Aman, M. G., & Gharabawi, G. M. (2004, September). Treatment of Behavior Disorders in Mental Retardation: Report on Transitioning to Atypical Antipsychotics, With an Emphasis on Risperidone. *The Journal of Clinical Psychiatry*, 65(9).
- American Diabetes Association, American Psychiatric Association, American Association of Clinical Endocrinologists, and North American Association for the Study of Obesity. (2004, February). Consensus Development Conference on Antipsychotic Drugs and Obesity and Diabetes. *Diabetes Care*, 27(2), 596-601.
- Amoako, E., Skelly, A. H., & Rossen, E. K. (2008, December). Outcomes of an Intervention to Reduce Uncertainty Among African American Women with Diabetes. *Western Journal of Nursing Research*, 30(8), 928-942.
- Andreescu, C., Lenze, E. J., Dew, M. A., Begley, A. E., Mulsant, B. H., Dombrowski, A. Y., et al. (2007, April). Effect of comorbid anxiety on treatment response and relapse risk in late-life depression: controlled study. *The British Journal of Psychiatry*, 190(4), 344-349.

Andros, V., Egger, A., & Dua, U. (2006, May). Blood Pressure Goal Attainment According to JNC 7 Guidelines and Utilization of Antihypertensive Drug Therapy in MCO Patients With Type 1 or Type 2 Diabetes. *Journal of Managed Care Pharmacy*, 12(4), 303-309.

Anonymous. (2000, March 11). Joint British recommendations on prevention of coronary heart disease in clinical practice: Summary. *British Medical Journal (International edition)*, 320(7236), 705-709.

Antochi, R., Stavrakaki, C., & Emergy, P. C. (2003, March). Psychopharmacological treatments in persons with dual diagnosis of psychiatric disorders and developmental disabilities. *Post Graduate Medical Journal*, 79(929), 139-146.

Anttila, S. K., Huhtala, H. S., Pekurinen, M. J., & Pitkääjärvi, T. K. (2000, March). Cost-effectiveness of an innovative four-year post-discharge programme for elderly patients. *Scandinavian Journal of Public Health*, 28(1), 41-47.

Arauz-Pacheco, C., Parrott, M. A., & Raskin, P. (2002, January). Position statement: Treatment of hypertension in adults with diabetes. *Diabetes Care*, 25(1), 199-202.

Austin, J., Williams, R., Ross, L., Moseley, L., & Hutchinson, S. (2005, March). Randomised controlled trial of cardiac rehabilitation in elderly patients with heart failure. *European Journal of Heart Failure*, 7(3), 411-417.

Ayalon, L., Feliciano, L., Gum, A. M., & Areán, P. A. (2006, November 13). Effectiveness of Nonpharmacological Interventions for the Management of Neuropsychiatric Symptoms in Patients With Dementia: A Systematic Review. *Archives of Internal Medicine*, 166(20), 2182-2188.

B

Babamoto, K. S., Sey, K. A., Camilleri, A. J., Karlan, V. J., Catalasan, J., & Moriksy, D. E. (2009, February). Improving Diabetes Care and Health Measures Among Hispanics Using Community Health Workers: Results From a Randomized Controlled Trial. *Health Education & Behavior*, 36(1), 113-126.

Bains, J., Birks, J., & Dening, T. (2002, October). Antidepressants for treating depression in dementia. *Cochrane Database of Systematic Reviews*(4).

Bao, Y., Post, E.P., Ten Have, T.R., Schackman, B.R., Bruce, M.L. (2009, May). Achieving Effective Antidepressant Pharmacotherapy in Primary Care: The Role of Depression Care Management in Treating Late-Life Depression. *Journal of the American Geriatrics Society*, 57(5), 895-900.

Barrowclough, C., Haddock, G., Tarrier, N., & Lewis, S. W. (2001, October). Randomized controlled trial of motivational interviewing, cognitive behavior therapy, and family intervention for patients with comorbid schizophrenia and substance use disorders. *The American Journal of Psychiatry*, 158(10), 1706-1714.

- Battersby, M., Harvey, P., Mills, P. D., Kalucy, E., Pols, R. G., Frith, P. A., et al. (2007). SA HealthPlus: A Controlled Trial of a Statewide Application of a Generic Model of Chronic Illness Care. *Milbank Quarterly*, 85(1), 37-67.
- Bauer, M. S., Biswas, K., & Kilbourne, A. M. (2009, November). Enhancing Multiyear Guideline Concordance for Bipolar Disorder Through Collaborative Care. *The American Journal of Psychiatry*, 166(11), 1244-1251.
- Bauer, M. S., McBride, L., Williford, W. O., & Glick, H. (2006, July). Collaborative Care for Bipolar Disorder: Part I. Intervention and Implementation in a Randomized Effectiveness Trial. *Psychiatric Services*, 57(7), 927.
- Bauer, M. S., McBride, L., Williford, W. O., & Glick, H. (2006, July). Collaborative Care for Bipolar Disorder: Part II. Impact on Clinical Outcome, Function, and Costs. *Psychiatric Services*, 57(7), 937.
- Bausewein, C., Booth, S., Gysels, M., & Higginson, I. (2008). Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases. *Cochrane Database of Systematic Reviews*(2).
- Bebb, C., Kendrick, D., Coupland, C., Madeley, R., Stewart, J., Brown, K., et al. (2007, February). A cluster randomised controlled trial of the effect of a treatment algorithm for hypertension in patients with type 2 diabetes. *The British Journal of General Practice*, 57(535), 136-143.
- Beesdo, K., Hartford, J., Russell, J., Spann, M., Ball, S., & Wittchen, H.U. (2009, December). The short- and long-term effect of duloxetine on painful physical symptoms in patients with generalized anxiety disorder: Results from three clinical trials. *Journal of Anxiety Disorders*, 23(8), 1064-1071.
- Belgrade, M. J., Cole, B. E., McCarberg, B. H., & McLean, M. J. (2006, April). Diabetic Peripheral Neuropathic Pain: Case Studies. *Mayo Clinic Proceedings*, 81(4), S12-25.
- Berthiaume J.T., Davis J., Taira D.A., & Thein K.K. (2007, March). A managed care organization's use of integrated health management to improve secondary prevention of coronary artery disease. *American Journal of Managed Care*, 13(3), 142-7.
- Beswick, A., Ebrahim, S., Burke, M., & Smith, G. (2009). Multiple risk factor interventions for primary prevention of coronary heart disease. *Cochrane Database of Systematic Reviews*, Issue 4.
- Betti, I., Castelli, G., Barchielli, A., Beligni, C., Boscherini, V., De Luca, L., et al. (2009, June). The Role of N-terminal PRO-Brain Natriuretic Peptide and Echocardiography for Screening Asymptomatic Left Ventricular Dysfunction in a Population at High Risk for Heart Failure. The PROBE-HF Study. *Journal of Cardiac Failure*, 15(5), 377-384.
- Bogner, H.R., and de Vries, H.F. (2008) Integration of Depression and Hypertension Treatment: A Pilot, Randomized Controlled Trial. *Annals of Family Medicine*, 6, 295-301.

- Bogner, H. R., Morales, K. H., Post, E. P., & Bruce, M. L. (2007, December). Diabetes, Depression, and Death: A randomized controlled trial of a depression treatment program for older adults based in primary care (PROSPECT). *Diabetes Care*, 30(12), 3005-3011.
- Bolen, S. D., Samuels, T. A., Yeh, H.C., Marinopoulos, S. S., McGuire, M., Abuid, M., et al. (2008, May). Failure to Intensify Antihypertensive Treatment by Primary Care Providers: A Cohort Study in Adults with Diabetes Mellitus and Hypertension. *Journal of General Internal Medicine*, 23(5), 543-550.
- Borzecki, A. M., & Berlowitz, D. R. (2005, December). Management of hypertension and diabetes: Review of treatment goals, drug choices, current practice and potential strategies for improvement in care. *Current Hypertension Reports*, 7(6), 439-449.
- Bosch, J., Glick, I. D., & Casey, D. E. (2009, June). A Double-Blind Randomized Trial of Mood Stabilizer Augmentation Using Lamotrigine and Valproate for Patients With Schizophrenia Who Are Stabilized and Partially Responsive. *Journal of Clinical Psychopharmacology*, 29(3), 267-271.
- Boult, C., Morishita, L., Dowd, B., Boult, L. B., Kane, R. L., & Urdangarin, C. F. (2001, April). A Randomized Clinical Trial of Outpatient Geriatric Evaluation and Management. *Journal of the American Geriatrics Society*, 49(4), 351-359.
- Boult, C., Reider, L., Frey, K., & Leff, B. (2008, March). Early Effects of "Guided Care" on the Quality of Health Care for Multimorbid Older Persons: A Cluster-Randomized Controlled Trial. *The Journals of Gerontology*, 63A(3), 321-327.
- Bourassa, M.G., & Berry, C. (2008, April). Prevention and noninvasive management of coronary atherosclerosis in patients with diabetes. *Current Atherosclerosis Reports*, 10(2), 106-16.
- Boyd, C. M., Boult, C., Shadmi, E., & Leff, B. (2007, October). Guided Care for Multimorbid Older Adults. *The Gerontologist*, 47(5), 697-705.
- Boyd, C. M., Shadmi, E., Conwell, L. J., Griswold, M., Leff, B., Brager, R., et al. (2008, May). A Pilot Test of the Effect of Guided Care on the Quality of Primary Care Experiences for Multimorbid Older Adults. *Journal of General Internal Medicine*, 23(5), 536-542.
- Briand, C., Vasiliadis, H.M., Lesage, A., Lalonde, P., Stip, E., Nicole, L., et al. (2006, July). Including Integrated Psychological Treatment as Part of Standard Medical Therapy for Patients With Schizophrenia: Clinical Outcomes. *Journal of Nervous & Mental Disease*, 194(7), 463-470.
- Brodsky, H., Draper, B. M., Millar, J., Low, L.F., Lie, D., Sharah, S., et al. (2003, January). Randomized Controlled Trial of Different Models of Care for Nursing Home Residents With Dementia Complicated by Depression or Psychosis. *The Journal of Clinical Psychiatry*, 64(1), 63-72.
- Brousseau, L., Wells, G.A., Finestone, H.M., Egan, M., Dubouloz, C.J., Graham, I., Casimiro, L., Robinson, V.A., Bilodeau, M., & McGowan, J. (2006, Spring). *Top Stroke Rehabilitation*, 13(2), 68-97.

Brown, E. L., Raue, P. J., & Halpert, K. D. (2007, June). Detecting depression in older adults with dementia. *University of Iowa Gerontological Nursing Interventions Research Center, Research Dissemination Core*, 39 p.

Brown, E. L., Raue, P., & Halpert, K. D. (2009). Evidence-Based Guideline: Detection of Depression in Older Adults with Dementia. *Journal of Gerontological Nursing*, 35(2), 11-15.

Brown, E.S., Murray, M, Carmody, T.J., Kennard, B.D., Hughes, C.W., Khan, D.A., & Rush, A.J. (2008, May). The Quick Inventory of Depressive Symptomatology-Self-report: a Psychometric Evaluation in Patients with Asthma and Major Depressive Disorder. *Department of Psychiatry, University of Texas Southwestern Medical Center*, 100(5), 433-438.

Brown, R. L., Saunders, L. A., Bobula, J. A., Mundit, M. P., & Koch, P. E. (2007, June). Randomized-Controlled Trial of a Telephone and Mail Intervention for Alcohol Use Disorders: Three-Month Drinking Outcomes. *Alcoholism: clinical and experimental research*, 31(8), 1372-1379.

Bruce M.L., Brown E.L., Raue P.J., Mlodzianowski A.E., Meyers B.S., Leon A.C., Heo M., Byers A.L., Greenberg R.L., Rinder S., Katt W., & Nassisi P. (2007, November). A randomized trial of depression assessment intervention in home health care. *Journal of the American Geriatric Society*, 55(11), 1793-1800.

Brunette, M. F., Noordsy, D. L., Xie, H., & Drake, R. E. (2003, October). Benzodiazepine Use and Abuse Among Patients With Severe Mental Illness and Co-occurring Substance Use Disorders. *Psychiatric Services*, 54(10), 1395-1401.

Bryson, C. L., Au, D. H., Sun, H., Williams, E. C., Kivlahan, D. R., & Bradley, K. A. (2008, December). Alcohol Screening Scores and Medication Nonadherence. *Annals of Internal Medicine*, 149(11), 795-803.

Burnam, M. A., & Watkins, K. E. (2006, May/June). Substance Abuse With Mental Disorders: Specialized Public Systems And Integrated Care. *Health Affairs*, 25(3), 648-658.

Butler, M., Kane, R. L., McAlpine, D., Kathol, R. G., Fu, S. S., Hagedorn, H., et al. (2008, November). Integration of mental health/substance abuse and primary care. *Evidence Report/Technology Assess (Full Rep)*, 173, 1-362.

C

Callahan, C. M., Boustani, M. A., Unverzagt, F. W., Austrom, M. G., Damush, T. M., Perkins, A. J., et al. (2006, May 10). Effectiveness of Collaborative Care for Older Adults With Alzheimer Disease in Primary Care: A Randomized Controlled Trial. *The Journal of the American Medical Association*, 295(18), 2148-2157.

Canzanello, V.J., Jensen, P.L., Schwartz, L.L., Worra, J.B., Klein, L.K. (2005). Improved Blood Pressure Control With a Physician-Nurse Team and Home Blood Pressure Measurement. *Mayo Clinic Proceedings*, 80(1), 31-36.

- Caplan G.A., Williams A.J., Daly B., & Abraham K. (2004, September). A randomized, controlled trial of comprehensive geriatric assessment and multidisciplinary intervention after discharge of elderly from the emergency department--the DEED II study. *Journal of the American Geriatric Society*, 52(9), 1417-1423.
- Capoccia, K. L., Boudreau, D. M., Blough, D. K., Ellsworth, A. J., Clark, D. R., Stevens, N. G., et al. (2004, February 15). Randomized trial of pharmacist interventions to improve depression care and outcomes in primary care. *American Journal of Health-System Pharmacy*, 61(4), 364-372.
- Care California Healthcare Foundation/American Geriatrics Society Panel in Improving Care for Elders with Diabetes. (2003, May). Guidelines for Improving the Care of the Older Person with Diabetes Mellitus. *Journal of the American Geriatrics Society*, 51(5S), 265-280.
- Carney, R.M., Freedland, K.E., Rubin, MD, E.H., Rich, M.W., Steinmeyer, B.C., Harris, W.S. (2009). Omega-3 Augmentation of Sertraline in Treatment of Depression in Patients With Coronary Heart Disease - A Randomized Controlled Trial. *The Journal of the American Medical Association*, 302(15), 1651-1657.
- Carter B.L., Rogers M., Daly J., Zheng S., James P.A. (2009, October). The potency of team-based care interventions for hypertension: a meta-analysis. *Archives of Internal Medicine*, 169(19), 1748-55.
- Casas, A., Troosters, T., Garcia-Aymerich, J., Roca, J., Hernández, C., Alonso, A., et al. (2006, July). Integrated care prevents hospitalisations for exacerbations in COPD patients. *European Respiratory Journal*, 28(1), 123-130.
- Chafetz, L., White, M., Collins-Bride, G., Nickens, J., & Cooper, B. A. (2008, June). Clinical Trial of Wellness Training: Health Promotion for Severely Mentally Ill Adults. *Journal of Nervous & Mental Disease*, 196(6), 475-483.
- Chan, J. C., So, W.Y., Yeung, C.Y., & Ko, G. T. (2009, June). Effects of Structured Versus Usual Care on Renal Endpoint in Type 2 Diabetes: The SURE Study: A randomized multicenter translational study. *Diabetes Care*, 32(6), 977-983.
- Chaput, Y., Magnan, A., & Gendron, A. (2008, August). The co-administration of quetiapine or placebo to cognitive-behavior therapy in treatment refractory depression: A preliminary trial. *BMC Psychiatry*, 8(73).
- Chen, W.C., Chu, H., Lu, R.-B., Chou, Y.H., Chen, C.H., Chang, Y.C., et al. (2009, August). Efficacy of progressive muscle relaxation training in reducing anxiety in patients with acute schizophrenia. *Journal of Clinical Nursing*, 18(15), 2187-2196.
- Chien, W. T., & Lee, Y. M. (2008, April). A Disease Management Program for Families of Persons in Hong Kong With Dementia. *Psychiatric Services*, 59(4), 433.
- Chobanian, A.V., Bakris, G.L., Black, H.R., Cushman, W.C., Green, L.A., Izzo, J.L., et al. (2003). The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment

of High Blood Pressure - The JNC 7 Report . *The Journal of the American Medical Association*, 289(19), 2560-2571.

Chodosh, J., Morton, S. C., Mojica, W., & Maglione, M. (2005, September 20). Meta-Analysis: Chronic Disease Self-Management Programs for Older Adults. *Annals of Internal Medicine*, 143(6), 427-438.

Choe, H. M., Cooke, D., Stutz, D., Standiford, C., & Berkstein, S. J. (2008, July-September). Using a Multidisciplinary Team and Clinical Redesign to Improve Blood Pressure Control in Patients With Diabetes. *Quality Management in Health Care*, 17(3), 227-233.

Clark, R. A., Inglis, S. C., McAlister, F. A., Cleland, J. G., & Stewart, S. (2007, April 10). Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. *BMJ*, 334(760).

Clark, R. E., Samnaliev, M., & McGovern, M. P. (2007, July). Treatment for Co-occurring Mental and Substance Use Disorders in Five State Medicaid Programs. *Psychiatric Services*, 58(7), 942-948.

Cleary, M., Hunt, G. E., Matheson, S. L., Siegfried, N., & Walter, G. (2008). Psychosocial interventions for people with both severe mental illness and substance misuse. *Cochrane Database of Systematic Reviews*(1).

Cleary, M., Hunt, G. E., Matheson, S. L., Siegfried, N., & Walter, G. (2008, January). Psychosocial interventions for people with both severe mental illness and substance misuse. *Cochrane Database of Systematic Reviews*, 23(1).

Coberley, C., Morrow, G., McGinnis, M., Wells, A., Coberley, S., Orr, P., Shurney, D. (2008, April). Increased Adherence to Cardiac Standards of Care during Participation in Cardiac Disease Management Programs. *Disease Management*, 11(2): 111-118.

Cohn, T. A., & Sernyak, M. J. (2006, July). Metabolic Monitoring for Patients Treated With Antipsychotic Medications. *Canadian Journal of Psychiatry*, 51(8), 492-501.

Cole, S. A., Farber, N. C., Weiner, J. S., Sulfano, M., Katzelnick, D. J., & Blader, J. C. (2006, October). Double-disease management or one care manager for two chronic conditions: pilot feasibility study of nurse telephonic disease management for depression and congestive heart failure. *Disease Management*, 9(5), 266-276.

Coleman, E.A., Parry, C., Chalmers, S., & Min, S.J. (2006, September). The care transitions intervention: results of a randomized controlled trial. *Archives of Internal Medicine*, 166(17), 1822-1828.

Collins, C., Levis Hewson, D., Munger, R., & Wade, T. (2010, May). Retrieved from <http://www.milbank.org/reports/10430EvolvingCare/10430EvolvingCare.html>.

Conley, R. R., Ascher-Svanum, H., Zhu, B., Faries, D. E., & Kinon, B. J. (2007, February). The burden of depressive symptoms in the long-term treatment of patients with schizophrenia. *Schizophrenia Research*, 90(1-3), 186-197.

Connor, K., McNeese-Smith, D., van Servellen, G., Chang, B., Lee, M., Cheng, E., et al. (2009, September-October). Insight into dementia care management using social-behavioral theory and mixed methods. *Nursing Research*, 58(5), 348-358.

Counsell, S. R., Callahan, C. M., Clark, D. O., Tu, W., Buttar, A. B., Stump, T. E., et al. (2007, December 12). Geriatric Care Management for Low-Income Seniors: A Randomized Controlled Trial. *The Journal of the American Medical Association*, 298(22), 2623-2633.

Coupar, F., Pollock, A., van Wijck, F., Morris, J., & Langhorne, P. (2010, April 14). Simultaneous bilateral training for improving arm function after stroke. *Cochrane Database of Systematic Reviews*(4).

Coventry, P. A., & Gellatly, J. L. (2008, September). Improving outcomes for COPD patients with mild-to-moderate anxiety and depression: A systematic review of cognitive behavioural therapy. *British Journal of Health Psychology*, 13(3), 381-400.

Coviello, D. M., Zanis, D. A., Wesnoski, S. A., & Alterman, A. I. (2006, October 15). The effectiveness of outreach case management in re-enrolling discharged methadone patients. *Drug and Alcohol Dependence*, 85(1), 56-65.

Craig, T. K., Johnson, S., McCrone, P., & Afuwape, S. (2008, March). Integrated Care for Co-occurring Disorders: Psychiatric Symptoms, Social Functioning, and Service Costs at 18 Months. *Psychiatric Services*, 59(3), 276-282.

Crotty, M., Halbert, J., Rowett, D., & Giles, L. (2004, November). An outreach geriatric medication advisory service in residential aged care: a randomised controlled trial of case conferencing. *Age and Ageing*, 33(6), 612-617.

Curran, G. M., Booth, B. M., Kirchner, J. E., & Deneke, D. E. (2007, August). Recognition and Management of Depression in a Substance Use Disorder Treatment Population. *American Journal of Drug & Alcohol Abuse*, 33(4), 563-569.

D

Daly, B., Abraham, K., Caplan, G. A., & Williams, A. J. (2004, September). A Randomized, Controlled Trial of Comprehensive Geriatric Assessment and Multidisciplinary Intervention After Discharge of Elderly from the Emergency Department—The DEED II Study. *Journal of the American Geriatrics Society*, 52(9), 1417-1423.

Damush, T. M., Wu, J., Bair, M. J., Sutherland, J. M., & Kroenke, K. (2008, August). Self-management practices among primary care patients with musculoskeletal pain and depression. *Journal of Behavioral Medicine*, 31(4), 301-307.

Davidson, K.W., Kupfer, D.J., Bigger, J.T., Califf, R.M., Carney, R.M., Coyne, J.C., Czajkowski, S.M., Frank, E., Frasure-Smith, N., Freedland, K.E., Froelicher, E.S., Glassman, A.H., et al. (2006, Sept-Oct). Assessment and treatment of depression in patients with cardiovascular disease: National Heart, Lung, and Blood Institute Working Group Report. *Psychosomatic Medicine*, 68(5), 645-650.

Davies, E. J., Moxham, T., Rees, K., Singh, S., Coats, A. J., Ebrahim, S., et al. (2010). Exercise based rehabilitation for heart failure. *Cochrane Database of Systematic Reviews*(4).

Davis, E., Saeed, S. A., & Antonacci, D. J. (2008, September). Anxiety Disorders in Persons with Developmental Disabilities: Empirically Informed Diagnosis and Treatment; Reviews Literature on Anxiety Disorders in DD Population with Practical Take-Home Messages for the Clinician. *Psychiatric Quarterly*, 79(3), 249-263.

Davis, L., Uezato, A., Newell, J.M., & Frazier, E. (2008, January). Major depression and comorbid substance use disorders. *Current Opinion in Psychology*, 21(1), 14-18.

Davison, S. N. (2007, December). The Prevalence and Management of Chronic Pain in End-Stage Renal Disease. *Journal of Palliative Medicine*, 10(6), 1277-1287.

de Jonge, P., Bel Hadj, F., Boffa, D., & Zdrojewski, C. (2009, May-June). Prevention of Major Depression in Complex Medically Ill Patients: Preliminary Results From a Randomized, Controlled Trial. *Psychosomatics*, 50(3), 227-233.

DeWalt, D. A., Malone, R. M., Bryant, M. E., Kosnar, M. C., Corr, K. E., Rothman, R. L., et al. (2006, March 13). A heart failure self-management program for patients of all literacy levels: A randomized, controlled trial. *BMC Health Services Research*, 6(30).

Diwan, S., & Phillips, V. L. (2001, March). Agitation and Dementia-Related Problem Behaviors and Case Management in Long-Term Care. *International Psychogeriatrics*, 13(1), 5-21.

Dobscha, S. K., Corson, K., Perrin, N. A., Hanson, G. C., Leibowitz, R. Q., Doak, M. N., et al. (2009, March). Collaborative Care for Chronic Pain in Primary Care: A Cluster Randomized Trial. *The Journal of the American Medical Association*, 301(12), 1242-1252.

Donald, M., Dower, J., & Kavanagh, D. (2005, March). Integrated versus non-integrated management and care for clients with co-occurring mental health and substance use disorders: a qualitative systematic review of randomised controlled trials. *Social Science & Medicine*, 60(6), 1371-1383.

Drake, R. E., O'Neal, E. L., & Wallach, M. A. (2008, January). A systematic review of psychosocial research on psychosocial interventions for people with co-occurring severe mental and substance use disorders. *Journal of Substance Abuse Treatment*, 34(1), 123-138.

Duggirala, M. K., Cuddihy, R. M., Cuddihy, M.T., Naessens, J. M., Cha, S. S., Mandrekar, J. N., et al. (2005, June). Predictors of Blood Pressure Control in Patients With Diabetes and Hypertension Seen in Primary Care Clinics. *American Journal of Hypertension*, 18(6), 833-838.

E

Eamonn Eeles, M. M. (2008, March). Delirium in the Long-Term Care Setting: Clinical and Research Challenges. *Journal of the American Medical Directors Association*, 9(3), 157-161.

Eiser, N., Harte, R., Karvounis, S., Phillips, C., & Isaac, M. T. (2005, June). Effect of Treating Depression on Quality-of-Life and Exercise Tolerance in Severe COPD. *COPD*, 2(2), 233-241.

Elkan, R., Kendrick, D., Dewey, M., & Hewitt, M. (2001, September 29). Effectiveness of home based support for older people: Systematic review and meta-analysis / Commentary. *British Medical Journal (International edition)*, 323(7315), 719-725.

Esposito, D., Brown, R., Chen, A., Schore, J., & Shapiro, R. (2008, Fall). Impacts of a Disease Management Program for Dually Eligible Beneficiaries. *Health Care Financing Review*, 30(1), 27-45.

F

Fellström, B. C., Jardine, A. G., Schmieder, R. E., & Holdaas, H. (2009, April 2). Rosuvastatin and Cardiovascular Events in Patients Undergoing Hemodialysis. *The New England Journal of Medicine*, 360(14), 1395-1407.

Ferro, J. M., & Pinto, F. (2004). Poststroke Epilepsy: Epidemiology, Pathophysiology and Management. *Drugs & Aging*, 21(10), 639-653.

Findling, R. L., Steiner, H., & Weller, E. B. (2005). Use of Antipsychotics in Children and Adolescents. *The Journal of Clinical Psychiatry*, 66(S7).

Flowers, L. (2007, March). Retrieved from AARP Public Policy Institute: assets.aarp.org/rgcenter/health/ib81_medicaid.pdf

Focht-New, G., Clements, P. T., Barol, B., Faulkner, M. J., & Pekala Service, K. (2008, January). Persons with Developmental Disabilities Exposed to Interpersonal Violence and Crime: Strategies and Guidance for Assessment. *Perspectives in Psychiatric Care*, 44(1), 3-13.

Forbes, D., Morgan, D. G., Markle-Reid, M., Wood, J., & Culum, I. (2008, July 16). Physical activity programs for persons with dementia. *Cochrane Database of Systematic Reviews*(3).

Fortney, J. C., Pyne, J. M., Edlund, M. J., Williams, D. K., Robinson, D. E., Mittal, D., et al. (2007, May 10). A Randomized Trial of Telemedicine-based Collaborative Care for Depression. *Journal of General Internal Medicine*, 22(8), 1086-1093.

Frangou, S., Sachpazidis, I., Stassinakis, A., & Sakas, G. (2005, December). Telemonitoring of medication adherence in patients with schizophrenia. *Telemedicine and e-Health*, 11(6), 675-683.

French, J.A. (2004, April 27). Efficacy and tolerability of the new antiepileptic drugs I: treatment of new onset epilepsy: report of the Therapeutics and Technology Assessment Subcommittee and Quality Standards Subcommittee of the American Academy of Neurology and the American Epileps. *Neurology*, 62(8), 1252-1260.

French, J.A. (2004, April 27). Efficacy and tolerability of the new antiepileptic drugs II: treatment of refractory epilepsy: report of the Therapeutics and Technology Assessment Subcommittee and Quality

Standards Subcommittee of the American Academy of Neurology and the American Epile. *Neurology*, 62(8), 1261-1273.

Friedman, B., Heisel, M., & Delavan, R. (2005, November). Validity of the SF-36 Five-Item Mental Health Index for Major Depression in Functionally Impaired, Community-Dwelling Elderly Patients. *Journal of the American Geriatrics Society*, 53(11), 1978-1985.

G

Gary, R. (2006, July). Exercise Self-Efficacy in Older Women with Diastolic Heart Failure: Results of a Walking Program and Education Intervention. *Journal of Gerontological Nursing*, 32(7), 31-39.

Gellis, Z. D., McClive-Reed, K. P., & Brown, E. L. (2009). Treatments for Depression in Older Persons with Dementia. *Annals of Long Term Care*, 17(2), 29-36.

Gensichen, J., von Korff, M., Peitz, M., Muth, C., Beyer, M., Güthlin, C., et al. (2009, September 15). Case Management for Depression by Health Care Assistants in Small Primary Care Practices - A Cluster Randomized Trial. *Annals of Internal Medicine*, 151(6), 369-378.

Gill, T. M., Baker, D. I., Gottschalk, M., & Peduzzi, P. (2002, October 3). A program to prevent functional decline in physically frail, elderly persons who live at home. *The New England Journal of Medicine*, 347(14), 1068-1074.

Giobbie-Hurder, A., Manheim, L., Henderson, W., Ulasevich, A., Cummings, J., Hughes, S. L., et al. (2000, December 13). Effectiveness of Team-Managed Home-Based Primary Care: A Randomized Multicenter Trial. *The Journal of the American Medical Association*, 284(22), 2877-2885.

Gitlin L.N., Winter L., Dennis M.P., Corcoran M., Schinfeld S., & Hauck W.W. (2006, May). A randomized trial of a multicomponent home intervention to reduce functional difficulties in older adults. *Journal of the American Geriatric Society*, 54(5), 809-816.

Gottlieb, S. S., Kop, W. J., Thomas, S. A., Katzen, S., Vesely, M. R., Greenberg, N., et al. (2007, May). A double-blind placebo-controlled pilot study of controlled-release paroxetine on depression and quality of life in chronic heart failure. *American Heart Journal*, 153(5), 868-873.

Graede, P., Vedel, P., Larsen, N., & Jensen, G. V. (2003, January 30). Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *The New England Journal of Medicine*, 348(5), 383-393.

Graham, H. (2004, December). Implementing integrated treatment for co-existing substance use and severe mental health problems in assertive outreach teams: training issues. *Drug & Alcohol Review*, 23(4), 463-470.

Green, A. I. (2007, January). Pharmacotherapy for schizophrenia and co-occurring substance use disorders. *Neurotoxicity Research*, 11(1), 33-39.

Greene, S. B., Reiter, K. L., Kilpatrick, K. E., Leatherman, S., Somers, S. A., & Hamblin, A. (2008, October-December). Searching for a business case for quality in Medicaid managed care. *Health Care Manage Rev*, 33(4), 350-360.

Gregg, E.W., & Brown, A. (2003, July). Cognitive and Physical Disabilities and Aging-Related Complications of Diabetes. *Clinical Diabetes*, 21(3), 113-118.

Griffiths, T. L., Burr, M. L., Campbell, I. A., & Lewis-Jenkins, V. (2000, January 29). Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: A randomised controlled trial. *The Lancet*, 355(9201), 362-368.

Grothues, J. M., Bischof, G., Reinhardt, S., Meyer, C., John, U., & Jürgen Rumpf, H. (2008, April 1). Effectiveness of brief alcohol interventions for general practice patients with problematic drinking behavior and comorbid anxiety or depressive disorders. *Drug and Alcohol Dependence*, 94(1-3), 214-220.

Gum, A. M., Areán, P. A., & Bostrom, A. (2007). Low-income depressed older adults with psychiatric comorbidity: secondary analyses of response to psychotherapy and case management. *International Journal of Geriatric Psychiatry*, 22(2), 124-130.

H

Hackett, M. L., Anderson, C. S., House, A., & Xia, J. (2008). Interventions for treating depression after stroke. *Cochrane Database of Systematic Reviews*(4).

Haddock, G., Barrowclough, C., Tarrier, N., Moring, J., O'Brien, R., Schofield, N., et al. (2003). Cognitive-behavioural therapy and motivational intervention for schizophrenia and substance misuse: 18-month outcomes of a randomised controlled trial. *The British Journal of Psychiatry*, 183(5), 418-426.

Hamilton, G., Aiken, L. S., Butner, J., Lockhart, C. A., Volk-Craft, B. E., & Williams, F. G. (2006, February). Outcome Evaluation of a Randomized Trial of the PhoenixCare Intervention: Program of Case Management and Coordinated Care for the Seriously Chronically Ill. *Journal of Palliative Medicine*, 9(1), 111-126.

Hansen, N. V., Jørgensen, T., & Ørtenblad, L. (2006). Massage and touch for dementia. *Cochrane Database of Systematic Reviews*(4).

Harpole, L. H., Williams, J. W., Olsen, M. K., Stechuchak, K. M., Oddone, E., Callahan, C. M., et al. (2005, January-February). Improving depression outcomes in older adults with comorbid medical illness. *General Hospital Psychiatry*, 27(1), 4-12.

Harris D, T. M. (2006). Caring for Australasians with Renal Impairment (CARI). *Nephrology (Carlton)*, 11 Suppl 1, S2-197.

Hebert, P. L., Sisk, J. E., Wang, J. J., Tuzzio, L., Casabianca, J. M., Chassin, M. R., et al. (2008, October 21). Cost-Effectiveness of Nurse-Led Disease Management for Heart Failure in an Ethnically Diverse Urban Community. *Annals of Internal Medicine*, 149(8), 540-548.

- Hesse, M., Vanderplasschen, W., Rapp, R., Broekaert, E., & Fridell, M. (2007, October 17). Case management for persons with substance use disorders. *Cochrane Database of Systematic Reviews*(4).
- Ho, P. M., Magid, D. J., Shetterly, S. M., Olson, K. L., Peterson, P. N., Masoudi, F. A., et al. (2008). Importance of Therapy Intensification and Medication Nonadherence for Blood Pressure Control in Patients With Coronary Disease. *Archives of Internal Medicine*, 168(3), 271-276.
- Hoerger, T.J., & Ahmann, A.J. (2008, February). The impact of diabetes and associated cardiometabolic risk factors on members: strategies for optimizing outcomes. *Journal of Managed Care Pharmacy*, 14(1 Suppl C), S2-14.
- Holsclaw, S.L., Olson, K.L., Hornak, R., & Denham, A.M. (2005, June). Assessment of patient satisfaction with telephone and mail interventions provided by a clinical pharmacy cardiac risk reduction service. *Journal of Managed Care Pharmacy*, 11(5), 403-409.
- Holtz-Eakin, D. (2004, October 13). An Analysis of the Literature on Disease Management Programs. Retrieved from <http://www.cbo.gov/doc.cfm?index=5909&type=0>
- Horsfall, J., Cleary, M., Walter, G., & Hunt, G. E. (2009, February). Psychosocial Treatments for People with Co-occurring Severe Mental Illnesses and Substance Use Disorders (Dual Diagnosis): A Review of Empirical Evidence. *Harvard Review of Psychiatry*, 17(1), 24-34.
- Howard, R., Ballard, C., O'Brien, J., & Burns, A. (2001, July). Guidelines for the management of agitation in dementia. *International Journal of Geriatric Psychiatry*, 16(7), 714-717.
- Howard, R., Sanders, R., & Lydall-Smith, S. M. (2008, August). The implementation of Restoring Health - a chronic disease model of care to decrease acute health care utilization. *Chronic Respiratory Disease*, 5(3), 133-141.
- Howlett, J. G., Mann, O. E., Bailie, R., Hathaway, R., Svendsen, A., Benoit, R., et al. (2009, September). Heart failure clinics are associated with clinical benefit in both tertiary and community care settings: Data from the Improving Cardiovascular Outcomes in Nova Scotia (ICONS) registry. *The Canadian Journal of Cardiology*, 25(9), e306-e311.
- Huang, T.T., Li, Y.T., & Wang, C.H. (2009, February). Individualized programme to promote self-care among older adults with asthma: randomized controlled trial. *Journal of Advanced Nursing*, 65(2), 348-358.
- Hudson, T. J., Owen, R. R., Thrush, C. R., Armitage, T. L., & Thapa, P. (2008, January). Guideline Implementation and Patient-Tailoring Strategies to Improve Medication Adherence for Schizophrenia. *Journal of Clinical Psychiatry*, 69(1), 74-80.
- Hughes, S.L., Weaver, F.M., Giobbie-Hurder, A., Manheim, L., Henderson, W., Kubal, J.D., Ulasevich, A., & Cummings, J. (2000). Effectiveness of Team-Managed Home-Based Primary Care - A Randomized Multicenter Trial. *The Journal of the American Medical Association*, 284, 2877-2885.

Hunt, S. A., Abraham, W.T., Chin, M. H., Feldman, A. M., Francis, G. S., Ganiats, T. G., et al. (2009, April 14). 2009 Focused Update Incorporated Into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines Developed. *Journal of the American College of Cardiology*, 53(15), e1-e90.

Huss, A., Stuck, A.E., Rubenstein, L.Z., Egger, M., & Clough-Gorr, K. M. (2008, March). Multidimensional Preventive Home Visit Programs for Community-Dwelling Older Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *The Journals of Gerontology*, 63(3), 298-307.

I

Ivanov, I., Klein, M., Green, W. H., & Coffey, B. (2006, December). The challenges of psychopharmacological management of children with severe developmental disabilities. *Journal of Child and Adolescent Psychopharmacology*, 16(6), 793-799.

J

Jaarsma, T., Lesman-Leegte, I., Luttik, M.-L., Hogenhuis, J., Sanderman, R., van der Wal, M. H., et al. (2008, February 11). Effect of Moderate or Intensive Disease Management Program on Outcome in Patients With Heart Failure: Coordinating Study Evaluating Outcomes of Advising and Counseling in Heart Failure (COACH). *Archives of Internal Medicine*, 168(3), 316-324.

Jabbour, A., Macdonald, P. S., Keogh, A. M., Kotlyar, E., Mellemkjaer, S., Coleman, C. F., et al. (2010, April 27). Differences Between Beta-Blockers in Patients With Chronic Heart Failure and Chronic Obstructive Pulmonary Disease: A Randomized Crossover Trial. *Journal of the American College of Cardiology*, 55(17), 1780-1787.

Jacob, R., & Chowdbury, A. (2008, January). Metabolic comorbidity in schizophrenia. *Indian Journal of Medical Sciences*, 62(1), 23-31.

Jansen, A. P., van Hout, H. P., van Marwijk, H. M., Nijpels, G., de Bruijne, M. C., Bosmans, J. E., et al. (2005, December 12). (Cost)-effectiveness of case-management by district nurses among primary informal caregivers of older adults with dementia symptoms and the older adults who receive informal care: design of a randomized controlled trial. *BMC Public Health*, 5(133).

Jarvis, C. M., Hayman, L. L., Braun, L. T., Schwertz, D. W., Ferrans, C. E., & Piano, M. R. (2007, November/December). Cardiovascular Risk Factors and Metabolic Syndrome in Alcohol- and Nicotine-Dependent Men and Women. *Journal of Cardiovascular Nursing*, 22(6), 429-435.

Jill Crandall, D. S. (2006, October). The Influence of Age on the Effects of Lifestyle Modification and Metformin in Prevention of Diabetes. *The Journals of Gerontology*, 61A(10), 1075-1081.

Jordan, N., Lee, T. A., Valenstein, M., & Weiss, K. B. (2007, October). Effect of Care Setting on Evidence-based Depression Treatment for Veterans with COPD and Comorbid Depression. *Journal of General Internal Medicine*, 22(10), 1447–1452.

K

Kalapatapu, R. K., & Neugroschl, J. A. (2009, April). Update on neuropsychiatric symptoms of dementia: Evaluation and management. *Geriatrics*, 64(4), 20-26.

Kane, R. L., Flood, S., Bershadsky, B., & Keckhafer, G. (2004, February). Effect of an Innovative Medicare Managed Care Program on the Quality of Care for Nursing Home Residents. *The Gerontologist*, 44(1), 95-103.

Kane, R. L., Homyak, P., Bershadsky, B., & Flood, S. (2006, July). Variations on a Theme Called PACE. *The Journals of Gerontology*, 61A(7), 689-693.

Kane, R. L., Homyak, P., Bershadsky, B., Lum, Y.-S., & Siadat, M. S. (2003, April). Outcomes of managed care of dually eligible older persons. *The Gerontologist*, 43(2), 165-174.

Kaner, E. F., Dickinson, H. O., Beyer, F. R., Campbell, F., Schlesinger, C., Heather, N., et al. (2007, April 18). Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database of Systematic Reviews*(2).

Karjalainen, K., Malmivaara, A., van Tulder, M., Roine, R., Jauhiainen, M., Hurri, H., et al. (2000, October). Biopsychosocial rehabilitation for upper limb repetitive strain injuries in working age adults. *Scandinavian Journal of Work, Environment & Health*, 26(5), 373-381.

Karlsson, M. R., Edner, M., Henriksson, P., Mejhert, M., Persson, H., Grut, M., et al. (2005, August). A nurse-based management program in heart failure patients affects females and persons with cognitive dysfunction most. *Patient Education and Counseling*, 58(2), 146-153.

Karp, J. F., Weiner, D. K., Dew, M. A., Begley, A., Miller, M. D., & Reynolds, C. F. (2010, June). Duloxetine and care management treatment of older adults with comorbid major depressive disorder and chronic low back pain: results of an open-label pilot study. *International Journal of Geriatric Psychiatry*, 25(6), 633-642.

Kaupila, L.I. (2009, June). Atherosclerosis and disc degeneration/low-back pain--a systematic review. *European Journal of Vascular and Endovascular Surgery*, 37(6), 661-670.

Katon, W., Unützer, J., Fan, M.Y., & Williams, J. W. (2006, February). Cost-Effectiveness and Net Benefit of Enhanced Treatment of Depression for Older Adults With Diabetes and Depression. *Diabetes Care*, 29(2), 265-270.

Katzelnick, D. J., Simon, G. E., Pearson, S. D., Manning, W. G., Helstad, C. P., Henk, H. J., et al. (2000, April). Randomized Trial of a Depression Management Program in High Utilizers of Medical Care. *Archives of Family Medicine*, 9(4), 345-351.

- Kemp, D. E., Gao, K., Ganocy, S. J., Caldes, E., Feldman, K., Chan, P. K., et al. (2009, July). Medical and substance use comorbidity in bipolar disorder. *Journal of Affective Disorders*, 116(1-2), 64-69.
- Kerr, E. A., Zikmund-Fisher, B. J., Kalmerus, M. L., Subramanian, U., Hogan, M. M., & Hofer, T. P. (2008, May 20). The role of clinical uncertainty in treatment decisions for diabetic patients with uncontrolled blood pressure. *Annals of Internal Medicine*, 148(10), 717-727.
- Khunti, K., Stone, M., Paul, S., Baines, J., Gisborne, L., Farooqi, A., et al. (2007, November). Disease management programme for secondary prevention of coronary heart disease and heart failure in primary care: a cluster randomised controlled trial. *Heart*, 93(11), 1398-1405.
- Kilbourne, A. M., Biswas, K., Pirraglia, P. A., Sajatovic, M., Williford, W. O., & Bauer, M. S. (2009, January). Is the collaborative chronic care model effective for patients with bipolar disorder and co-occurring conditions? *Journal of Affective Disorders*, 112(1-3), 256-261.
- Kleber H.D., Weiss R.D., Anton R.F. Jr, George T.P., Greenfield S.F., Kosten T.R., O'Brien C.P., et al. (2007, April). Treatment of patients with substance use disorders, second edition. *American Journal of Psychiatry*, 164(4 Suppl), 5-123.
- Kolanowski, A., Fick, D., Waller, J. L., & Ahern, F. (2006, October). Outcomes of Antipsychotic Drug Use in Community-Dwelling Elders With Dementia. *Archives of Psychiatric Nursing*, 20(5), 217-225.
- Kotlerman, J.B., Afifi, A.A., Morisky, D.E., & Kominski, G.F. (2007, June). Impact of disease management on health care utilization: Evidence from the "Florida: A Healthy State (FAHS)" Medicaid Program. *Preventative Medicine*, 44(6), 547-553.
- Kotlyar, E., Keogh, A.M., Macdonald, P.S., Arnold, R.H., McCaffrey, D. J., & Glanville, A. R. (2002, December). Tolerability of carvedilol in patients with heart failure and concomitant chronic obstructive pulmonary disease or asthma. *The Journal of Heart and Lung Transplantation*, 21(12), 1290-1295.
- Kraus, C. A., Seignourel, P., Balasubramanyam, V., Snow, A.L., Wilson, N.L., Kunik, M.E., et al. (2008, May). Cognitive-behavioral treatment for anxiety in patients with dementia: two case studies. *Journal of Psychiatric Practice*, 14(3), 186-192.
- Kroenke, K., Krebs, E.E., & Bair, M.J. (2009, May-June). Pharmacotherapy of chronic pain: a synthesis of recommendations from systematic reviews. *General Hospital Psychiatry*, 31(3), 206-219.
- Kroenke, K., Wu, J., Hoke, S., Sutherland, J., Tu, W., Bair, M.J., et al. (2009, May 27). Optimized Antidepressant Therapy and Pain Self-management in Primary Care Patients With Depression and Musculoskeletal Pain: A Randomized Controlled Trial. *The Journal of the American Medical Association*, 301(20), 2099-2110.
- Kunik, M.E., Azzam, P.N., Soucek, J., & Cully, J.A. (2007, January-February). A Practical Screening Tool for Anxiety and Depression in Patients With Chronic Breathing Disorders. *Psychosomatics*, 48(1), 16-21.

Kunik, M.E., Veazey, C., Cully, J.A., & Soucek, J. (2008, March). COPD education and cognitive behavioral therapy group treatment for clinically significant symptoms of depression and anxiety in COPD patients: a randomized controlled trial. *Psychological Medicine*, 38(3), 385-396.

Kwan, J., & Wood, E. (2010, January 20). Antiepileptic drugs for the primary and secondary prevention of seizures after stroke. *Cochrane Database of Systematic Reviews*(1).

L

Lacasse, Y., Goldstein, R., Lasserson, T.J., & Martin, S. (2006, June 16). Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*(4).

Landis, S.E., Gaynes, B.N., Morrissey, J.P., Vinson, N., Ellis, A.R., & Domino, M.E. (2007, March 5). Generalist care managers for the treatment of depressed medicaid patients in North Carolina: A pilot study. *BMC Family Practice*, 8(7).

Lane, D. A., Chong, A. Y., & Lip, G. Y. (2005, January 24). Psychological interventions for depression in heart failure. *Cochrane Database of Systematic Reviews*(1).

Lang, A. J., Norman, G. J., & Casmar, P. V. (2006, December). A randomized trial of a brief mental health intervention for primary care patients. *Journal of Consulting and Clinical Psychology*, 74(6), 1173-1179.

Le Jemtel, T. H., Padeletti, M., & Jelic, S. (2007, January 16). Diagnostic and Therapeutic Challenges in Patients With Coexistent Chronic Obstructive Pulmonary Disease and Chronic Heart Failure. *Journal of the American College of Cardiology*, 49(2), 171-180.

Leff, B., Burton, L., Mader, S. L., & Naughton, B. (2005, December 6). Hospital at Home: Feasibility and Outcomes of a Program To Provide Hospital-Level Care at Home for Acutely Ill Older Patients. *Annals of Internal Medicine*, 143(11), 798-808.

Leff, B., Reider, L., Frick, K. D., Scharfstein, D. O., Boyd, C. M., Frey, K., et al. (2009, August). Guided care and the cost of complex healthcare: a preliminary report. *The American Journal of Managed Care*, 15(8), 555-559.

Lehrer, P. M., Katsamanis Karavidas, M., Lu, S.E., Feldman, J., Kranitz, L., Abraham, S., et al. (2008, May). Psychological treatment of comorbid asthma and panic disorder: A pilot study. *Journal of Anxiety Disorders*, 22(4), 671-683.

Leichsenring, F., Salzer, S., Jaeger, U., & Kächele, H. (2009, August). Short-Term Psychodynamic Psychotherapy and Cognitive-Behavioral Therapy in Generalized Anxiety Disorder: A Randomized, Controlled Trial. *The American Journal of Psychiatry*, 166(8), 875-881.

Lenze, E. J., Mulsant, B. H., Dew, M. A., Shear, M. K., Houck, P., Pollock, B. G., et al. (2003, December). Good treatment outcomes in late-life depression with comorbid anxiety. *Journal of Affective Disorders*, 77(3), 247-254.

- Lewis, A. (2004, Spring). Savings opportunities through Medicaid disease management. *Disease Management*, 7(1), 35-46.
- Lin, E. H., Tang, L., Katon, W., Hegel, M. T., Sullivan, M. D., & Unützer, J. (2006, November-December). Arthritis pain and disability: response to collaborative depression care. *General Hospital Psychiatry*, 28(6), 482-486.
- Lind, A., & Kaplan, L. (2007, October). An evaluation of Washington's Medicaid disease-management program. *Managed Care Interface*, 20(10), 45-50.
- Linden, A., Berg, G. D., & Wadhwa, S. (2007, October). Evaluation of a medicaid asthma disease management program. *Disease Management*, 10(5), 266-272.
- Lolak, S., Connors, G. L., Sheridan, M. J., & Wise, T. N. (2008, January). Effects of Progressive Muscle Relaxation Training on Anxiety and Depression in Patients Enrolled in an Outpatient Pulmonary Rehabilitation Program. *Psychotherapy and Psychosomatics*, 77(2), 119-125.
- Lucie Brosseau PhD, G. A.J. (2006, Spring). Ottawa Panel Evidence-Based Clinical Practice Guidelines for Post-Stroke Rehabilitation. *Topics in Stroke Rehabilitation*, 13(2), 1-269.
- Lustman, P. J., Freedland, K. E., Griffith, L. S., & Clouse, R. E. (2000, May). Fluoxetine for depression in diabetes: A randomized double-blind placebo-controlled trial. *Diabetes Care*, 23(5), 618-623.
- Lyketsos, C. G., Colenda, C. C., Beck, C., & Blank, K. (2006, July). Position Statement of the American Association for Geriatric Psychiatry Regarding Principles of Care for Patients With Dementia Resulting From Alzheimer Disease. *The American Journal of Geriatric Psychiatry*, 14(7), 561-572.
- ## M
- Maltais, F., Bourbeau, J., Shapiro, S., Lacasse, Y., Perrault, H., Baltzan, M., et al. (2008, December 16). Effects of Home-Based Pulmonary Rehabilitation in Patients with Chronic Obstructive Pulmonary Disease. *Annals of Internal Medicine*, 49(12), 869-878.
- Mancia, G. (2007, June). Optimal control of blood pressure in patients with diabetes reduces the incidence of macro and microvascular events. *Journal of Hypertension. Supplement: Official Journal of the International Society of Hypertension*, 25(1), S7-12.
- Manu, P., Kane, J. M., Frederickson, A. M., & Correll, C. U. (2008, November). Equally increased risk for metabolic syndrome in patients with bipolar disorder and schizophrenia treated with second-generation antipsychotics. *Bipolar Disorders*, 10(7), 788-797.
- Markle-Reid, M., Browne, G., Weir, R., Gafni, A., Roberts, J., & Henderson, S. R. (2006, October). The Effectiveness and Efficiency of Home-Based Nursing Health Promotion for Older People: A Review of the Literature. *Medical Care Research and Review*, 63(5), 531-569.
- Martin, D. C., Berger, M. L., Anstatt, D. T., Turpin, R. S., Cannuscio, C. C., Teutsch, S. M., et al. (2004, October). A Randomized Controlled Open Trial of Population-based Disease and Case

- Management in a Medicare Plus Choice Health Maintenance Organization. *Preventing Chronic Disease*, 1(4), A05.
- Mascarenhas, J., Lourenço, P., Lopes, R., Azevedo, A., & Bettencourt, P. (2008, March). Chronic obstructive pulmonary disease in heart failure. Prevalence, therapeutic and prognostic implications. *American Heart Journal*, 155(3), 521-525.
- Masi, G. (2004). Pharmacotherapy of Pervasive Developmental Disorders in Children and Adolescents. *CNS Drugs*, 18(14), 1031-1052.
- McAlister, F. A., Stewart, S., Ferrua, S., & McMurray, J. J. (2004, August 18). Multidisciplinary strategies for the management of heart failure patients at high risk for admission: A systematic review of randomized trials. *Journal of the American College of Cardiology*, 44(4), 810-819.
- McConnell, K.J., Humphries, T.L., Raebel, M.A., & Merenich, J.A. (2003, December). Clinical pharmacy specialist implementation of lisinopril therapy in patients with coronary artery disease and diabetes mellitus. *Pharmacotherapy*, 23(12), 1564-1572.
- McCord, J., Jneid, H., Hollander, J. E., de Lemos, J. A., Cercek, B., Hsue, P., et al. (2008, April 8). Management of cocaine-associated chest pain and myocardial infarction: a scientific statement from the American Heart Association Acute Cardiac Care Committee of the Council on Clinical Cardiology. *Circulation*, 117(14), 1897-1907.
- McCrone, P., Killaspy, H., Bebbington, P., Johnson, S., Nolan, F., Pilling, S., et al. (2009, Jul). The REACT study: cost-effectiveness analysis of assertive community treatment in north London. *Psychiatric Services*, 60(7), 908-913.
- McDermott, S., Moran, R., Platt, T., & Dasari, S. (2006, June). Variation in health conditions among groups of adults with disabilities in primary care. *Journal of Community Health*, 31(3), 147-159.
- McDermott, S., Moran, R., Platt, T., & Dasari, S. (2007, June). Health conditions among women with a disability. *Journal of Women's Health*, 16(5), 713-720.
- McGurk, S. R., Mueser, K. T., Derosa, T. J., & Wolfe, R. (2009, March). Work, Recovery, and Comorbidity in Schizophrenia: A Randomized Controlled Trial of Cognitive Remediation. *Schizophrenia Bulletin*, 35(2), 319-335.
- McKay, J. R., Lynch, K. G., Shepard, D. S., & Pettinati, H. M. (2005, February). The Effectiveness of Telephone-Based Continuing Care for Alcohol and Cocaine Dependence: 24-Month Outcomes. *The Archives of General Psychiatry*, 62(2), 199-207.
- McLean, D. L., McAlister, F. A., Johnson, J. A., King, K. M., Makowsky, M. J., Jones, C. A., et al. (2008, November 24). A randomized trial of the effect of community pharmacist and nurse care on improving blood pressure management in patients with diabetes mellitus: study of cardiovascular risk intervention by pharmacists-hypertension (SCRIP-HTN). *Archives of Internal Medicine*, 168(21), 2355-2361.

- McQueen, J., Howe, T. E., Allan, L., & Mains, D. (2009, July 8). Brief interventions for heavy alcohol users admitted to general hospital wards. *Cochrane Database of Systematic Reviews*(3).
- Mertens, J. R., Flisher, A. J., Satre, D. D., & Weisner, C. M. (2008, November 1). The role of medical conditions and primary care services in 5-year substance use outcomes among chemical dependency treatment patients. *Drug and Alcohol Dependence*, 98(1-2), 45-53.
- Meulepas, M. A., Jacobs, J. E., Smeenk, F. W., Smeele, I., Lucas, A. E., Bottema, B. J., et al. (2007, September). Effect of an integrated primary care model on the management of middle-aged and old patients with obstructive lung diseases. *Scandinavian Journal of Primary Health Care*, 25(3), 186-192.
- Milani, R.V., Lavie, C.J. (2007, September). Impact of Cardiac Rehabilitation on Depression and Its Associated Mortality. *The American Journal of Medicine*, 120(9).
- Millar, H. (2008, May). Management of physical health in schizophrenia: a stepping stone to treatment success. *European Neuropsychopharmacology*, 18(Suppl 2), S121-128.
- Minkoff, K. (2001, May). Best Practices: Developing Standards of Care for Individuals With Co-occurring Psychiatric and Substance Use Disorders. *Psychiatric Services*, 52, 597-599.
- Monette, J., Champoux, N., Monette, M., Fournier, L., Wolfson, C., du Fort, G. G., et al. (2008, June). Effect of an interdisciplinary educational program on antipsychotic prescribing among nursing home residents with dementia. *International Journal of Geriatric Psychiatry*, 23(6), 574-579.
- Morken, G., Grawe, R. W., & Widen, J. H. (2007, April). Effects of integrated treatment on antipsychotic medication adherence in a randomized trial in recent-onset schizophrenia. *The Journal of Clinical Psychiatry*, 68(4), 566-571.
- Murray, M. D., Ritchey, M. E., Wu, J., & Tu, W. (2009, April 27). Effect of a pharmacist on adverse drug events and medication errors in outpatients with cardiovascular disease. *Archives of Internal Medicine*, 169(8), 757-763.
- Myint, P. K., Staufenberg, E. F., & Sabanathan, K. (2006, September). Post-stroke seizure and post-stroke epilepsy. *Post Graduate Medical Journal*, 82(971), 568-572.

N

- Nasrallah, H. A., Meyer, J. M., Goff, D. C., McEvoy, J. P., Davis, S. M., Stroup, T. S., et al. (2006, September). Low rates of treatment for hypertension, dyslipidemia and diabetes in schizophrenia: data from the CATIE schizophrenia trial sample at baseline. *Schizophrenia Research*, 86(1-3), 15-22.
- National Kidney Foundation. (2007). KDOQI Clinical Practice Guidelines and Clinical Practice Recommendations for Diabetes and Chronic Kidney Disease. Retrieved from http://www.kidney.org/professionals/kdoqi/guideline_diabetes/
- National Stroke Foundation. (2005, September 8). Clinical Guidelines for Stroke Rehabilitation and Recovery. *National Stroke Foundation*. Melbourne, Australia.

Nguyen, V., Ducharme, A., White, M., Racine, N., O'Meara, E., Zhang, B., et al. (2007, May). Lack of long-term benefits of a 6-month heart failure disease management program. *Journal of Cardiac Failure*, 13(4), 287-293.

Niesink, A. (2007, November). Systematic review of the effects of chronic disease management on quality-of-life in people with chronic obstructive pulmonary disease. *Respiratory Medicine*, 101(11), 2233-2239.

Noble, M. J., Smith, J. R., & Windley, J. (2006, April). A controlled retrospective pilot study of an 'at-risk asthma register' in primary care. *Primary Care Respiratory Journal*, 15(2), 116-124.

Norris C.M., Patterson L., Galbraith D., Hegadoren K.M. (2009, May). All you have to do is call; a pilot study to improve the outcomes of patients with coronary artery disease. *Applied Nursing Research*, 22(2), 133-137.

O

Ohayon, M. M., & Schatzberg, A. F. (2003, January). Using chronic pain to predict depressive morbidity in the general population. *Archives of General Psychiatry*, 60(1), 39-47.

O'Kane, M. J., Bunting, B., Copeland, M., & Coates, V. E. (2008, May). Efficacy of self monitoring of blood glucose in patients with newly diagnosed type 2 diabetes (ESMON study): randomised controlled trial. *BMJ*, 336(7654), 1174-1177.

Orozco, L. J., Buchleitner, A. M., Gimenez-Perez, G., Roqué i Figuls, M., Richter, B., & Mauricio, D. (2008, July 16). Exercise or exercise and diet for preventing type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews*(3).

Oslin, D. W., Sayers, S., Ross, J., Kane, V., Ten Have, T., Conigliaro, J., et al. (2003, November-December). Disease management for depression and at-risk drinking via telephone in an older population of veterans. *Psychosomatic Medicine*, 65(6), 931-937.

P

Papademetriou V., Papadopoulos DP. (2008 May). Aggressive blood pressure control and stroke prevention: role of calcium channel blockers. *Journal of Hypertension*, 26(5), 844-852.

Pariser, D., & O'Hanlon, A. (2005). Effects of telephone intervention on arthritis self-efficacy, depression, pain, and fatigue in older adults with arthritis. *Journal of Geriatric Physical Therapy*, 28(3), 68-73.

Parry, C., Chalmers, S., Min, S.J., & Coleman, E. A. (2006, September 25). The Care Transitions Intervention: Results of a Randomized Controlled Trial. *Archives of Internal Medicine*, 166(17), 1822-1828.

- Passmore, M. J., Gardner, D. M., Polak, Y., & Rabheru, K. (2008). Alternatives to atypical antipsychotics for the management of dementia-related agitation. *Drugs & Aging*, 25(5), 381-398.
- Patel, R. R., Saltoun, C. A., & Grammer, L. C. (2009, February). Improving asthma care for the elderly: a randomized controlled trial using a simple telephone intervention. *Journal of Asthma*, 46(1), 30-35.
- Peikes, D., Chen, A., Schore, J., & Brown, R. (2009, February 11). Effects of care coordination on hospitalization, quality of care, and health care expenditures among Medicare beneficiaries: 15 randomized trials. *The Journal of the American Medical Association*, 301(6), 603-618.
- Pellegrini, F., Belfiglio, M., De Berardis, G., Franciosi, M., Di Nardo, B., Greenfield, S., et al. (2003, February 24). Role of organizational factors in poor blood pressure control in patients with type 2 diabetes: the QuED Study Group--quality of care and outcomes in type 2 diabetes. *Archives of Internal Medicine*, 163(4), 473-480.
- Perry, E., Swartz, J., Brown, S., Smith, D., Kelly, G., & Swartz, R. (2005, July). Peer mentoring: a culturally sensitive approach to end-of-life planning for long-term dialysis patients. *American Journal of Kidney Diseases*, 46(1), 111-119.
- Peterson, L. A., Woodard, L. D., Henderson, L. M., Urech, T. H., & Pietz, K. (2009, June 16). Will hypertension performance measures used for pay-for-performance programs penalize those who care for medically complex patients? *Circulation*, 119(23), 2978-2985.
- Petrakis, I. L., O'Malley, S., Rounsaville, B., Poling, J., McHugh-Strong, C., & Krystal, J. H. (2004, March). Naltrexone augmentation of neuroleptic treatment in alcohol abusing patients with schizophrenia. *Psychopharmacology (Berl)*, 172(3), 291-297.
- Phibbs, C. S., Holty, J. E., Goldstein, M. K., Garber, M. K., Wang, Y., Feussner, J. R., et al. (2006, January). The effect of geriatrics evaluation and management on nursing home use and health care costs: results from a randomized trial. *Medical Care*, 44(1), 91-95.
- Phillips, C. O., Wright, S. M., Kern, D. E., Shepperd, S., Singa, R. M., & Rubin, H. R. (2004, March 17). Comprehensive Discharge Planning With Postdischarge Support for Older Patients With Congestive Heart Failure: A Meta-analysis. *The Journal of the American Medical Association*, 291(11), 1358-1367.
- Pladeck, T., Hader, C., Von Orde, A., Rasche, K., & Wiechmann, H. W. (2007, November). Non-invasive ventilation: comparison of effectiveness, safety, and management in acute heart failure syndromes and acute exacerbations of chronic obstructive pulmonary disease. *The Journal of Physiology and Pharmacology*, 58(Supplement 5 (part 2)), 539-549.
- Planas, L. G., Crosby, K. M., Mitchell, K. D., & Farmer, K. C. (2009, March-April). Evaluation of a hypertension medication therapy management program in patients with diabetes. *Journal of the American Pharmacists Association*, 49(2), 164-170.

Post, K., Nutting, P., Smith, J. L., Elliott, C. E., & Dickinson, M. (2002, October 26). Managing depression as a chronic disease: A randomised trial of ongoing treatment in primary care. *British Medical Journal (International edition)*, 325(7370), 934-937.

Powell, H., & Gibson, P. G. (2003, January 20). Options for self-management education for adults with asthma. *Cochrane Database of Systematic Reviews*(1).

Powell, J., Chiu, T., & Eysenbach, G. (2008). A systematic review of networked technologies supporting carers of people with dementia. *Journal of Telemedicine and Telecare*, 14(3), 154-156.

Praet, S. F., van Rooij, E. S., Wijtvliet, A., & Boonman-deWinter, L. J. (2008, May). Brisk walking compared with an individualised medical fitness programme for patients with type 2 diabetes: a randomised controlled trial. *Diabetologia*, 51(5), 736-746.

Price, L. M. (2007, December). Transition to Community: A Program to Help Clients with Schizophrenia Move from Inpatient to Community Care; A Pilot Study. *Archives of Psychiatric Nursing*, 21(6), 336-344.

Pugh, M. J., Foreman, P. J., & Berlowitz, D. R. (2006). Prescribing antiepileptics for the elderly: differences between guideline recommendations and clinical practice. *Drugs & Aging*, 23(11), 861-875.

Pugh, M. J., Vancott, A. C., Steinman, M. A., Mortensen, E. M., Amuan, M. E., Wang, C. P., et al. (2010, March). Choice of initial antiepileptic drug for older veterans: possible pharmacokinetic drug interactions with existing medications. *Journal of the American Geriatrics Society*, 58(3), 465-471.

Puhan, M., Scharplatz, M., Troosters, T., Walters, E. H., & Steurer, J. (2009). Pulmonary rehabilitation following exacerbations of chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*(1).

R

Rabindranath, K. S., Butler, J., Roderick, P. J., Wallace, S. A., Daly, C., & MacLeod, A. M. (2005). Physical measures for treating depression in dialysis patients. *Cochrane Database of Systematic Reviews*(2).

Rabow, M. W., Dibble, S. L., Pantilat, S. Z., & McPhee, S. J. (2004, January 12). The Comprehensive Care Team: A Controlled Trial of Outpatient Palliative Medicine Consultation. *Archives of Internal Medicine*, 164(1), 83-91.

Rachmani, R., Levi, Z., Slavachevski, I., Avin, M., & Ravid, M. (2002, May). Teaching patients to monitor their risk factors retards the progression of vascular complications in high-risk patients with Type 2 diabetes mellitus--a randomized prospective study. *Diabetic Medicine*, 19(5), 385-392.

Reboldi, G., Gentile, G., Angeli, F., & Verdecchia, P. (2009, November). Exploring the optimal combination therapy in hypertensive patients with diabetes mellitus. *Expert Review of Cardiovascular Therapy*, 7(11), 1349-1361.

Riegel, B., Carlson, B., Glasner, D., & Romero, T. (2006, April). Randomized Controlled Trial of Telephone Case Management in Hispanics of Mexican Origin With Heart Failure. *Journal of Cardiac Failure*, 12(3), 211-219.

Rodríguez-Morán, M., Barragán-Rodríguez, L., & Guerrero-Romero, F. (2008, December). Efficacy and safety of oral magnesium supplementation in the treatment of depression in the elderly with type 2 diabetes: a randomized, equivalent trial. *Magnesium Research*, 21(4), 218-223.

Rose, H. L., Miller, P. M., Nemeth, L. S., Jenkins, R. G., Nietert, P. J., Wessell, A. M., et al. (2008, August). Alcohol screening and brief counseling in a primary care hypertensive population: a quality improvement intervention. *Addiction*, 103(8), 1271-1280.

Rose, L., & Gerdtz, M. F. (2009, December). Review of non-invasive ventilation in the emergency department: clinical considerations and management priorities. *Journal of Clinical Nursing*, 18(23), 3216-3224.

Rosenzweig, J.L., Ferrannini, E., Grundy, S.M., Haffner, S.M., Heine, R.J., Horton, E.S., & Kawamori, R. (2008). Primary Prevention of Cardiovascular Disease and Type 2 Diabetes in Patients at Metabolic Risk: An Endocrine Society Clinical Practice Guideline. *The Journal of Clinical Endocrinology & Metabolism*, 93(10), 3671-3689.

Royal Australian and New Zealand College of Psychiatrists clinical practice guidelines for the treatment of schizophrenia and related disorders. (2005, January-February). *The Australian and New Zealand Journal of Psychiatry*, 39(1-2), 1-30.

Rutten, F. H., Moons, K. G., Cramer, M.-J. M., Grobbee, D. E., Zuithoff, N. P., Lammers, J.W., et al. (2005, December 10). Recognising Heart Failure In Elderly Patients With Stable Chronic Obstructive Pulmonary Disease In Primary Care: Cross Sectional Diagnostic Study. *British Medical Journal*, 331(7529), 1379-1382.

Rydén L., Standl E., Bartnik M., Van den Berghe G., Betteridge J., de Boer M.J., Cosentino F., Jönsson B., Laakso M., et al. (2007, January). Guidelines on diabetes, pre-diabetes, and cardiovascular diseases: executive summary. The Task Force on Diabetes and Cardiovascular Diseases of the European Society of Cardiology (ESC) and of the European Association for the Study of Diabetes (EASD). *European Heart Journal*, 28(1), 88-136.

S

Sabaawi, M., Singh, N. N., & de Leon, J. (2006, May-June). Guidelines for the use of clozapine in individuals with developmental disabilities. *Research in Developmental Disabilities*, 27(3), 309-336.

Saenz, A., Fernandez-Esteban, I., Mataix, A., Ausejo Segura, M., Roqué i Figuls, M., & Moher, D. (2005, July 20). Metformin monotherapy for type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews*(3).

- Safaei, N. (2008, November 15). Outcomes of coronary artery bypass grafting in patients with a history of opiate use. *Pakistan Journal of Biological Sciences*, 11(22), 2594-2598.
- Salisbury, E. M., Game, D. S., Al-Shakarchi, I., Chan, M., Fishman, L., Tookman, L., et al. (2009, January). Changing practice to improve pain control for renal patients. *Postgraduate Medical Journal*, 85(999), 30-33.
- Salpeter, S. R., Ormiston, T. M., & Salpeter, E. E. (2005). Cardioselective beta-blockers for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*(4).
- San, L., Arranz, B., & Martinez-Raga, J. (2007). Antipsychotic drug treatment of schizophrenic patients with substance abuse disorders. *European Addiction Research*, 13(4), 230-243.
- Sandhoff, B.G., Nies, L.K., Olson, K.L., Nash, J.D., Rasmussen, J.R., & Merenich J.A. (2007). Clinical pharmacy cardiac risk service for managing patients with coronary artery disease in a health maintenance organization. *American Journal of Health-System Pharmacy*, 64(1), 77-84.
- Sanders, K. M., & Satyvavolu, A. (2002, Winter). Improving blood pressure control in diabetes: limitations of a clinical reminder in influencing physician behavior. *The Journal of Continuing Education in the Health Professions*, 22(1), 23-32.
- Schatz, M., Gibbons, C., Nelle, C., Harden, K., & Zeiger, R. S. (2006, April). Impact of a care manager on the outcomes of higher risk asthmatic patients: a randomized controlled trial. *The Journal of Asthma*, 43(3), 225-229.
- Scherr, D., Kastner, P., Kollman, A., Hallas, A., Auer, J., Krappinger, H., et al. (2009, August 17). Effect of home-based telemonitoring using mobile phone technology on the outcome of heart failure patients after an episode of acute decompensation: randomized controlled trial. *Journal of Medical Internet Research*, 11(3), e34.
- Schneider, L. S., Dagerman, K. S., & Insel, P. (2005, October 19). Risk of Death With Atypical Antipsychotic Drug Treatment for Dementia: Meta-analysis of Randomized Placebo-Controlled Trials. *The Journal of the American Medical Association*, 294(15), 1934-1943.
- Schneiderhan, M.E., Batscha, C.L., & Rosen, C. (2009, August). Assessment of a point-of-care metabolic risk screening program in outpatients receiving antipsychotic agents. *Pharmacotherapy*, 29(8), 975-987.
- Schoenthaler, A., Ogedegbe, G., Allegrante, J.P. (2009, February). Self-efficacy mediates the relationship between depressive symptoms and medication adherence among hypertensive African Americans. *Health Education & Behavior*, 36(1), 127-37.
- Seignourel, P. J., Kunik, M. E., Snow, L., Wilson, N., & Stanley, M. (2008, October). Anxiety in dementia: a critical review. *Clinical Psychology Review*, 28(7), 1071-1082.

- Sheehan, D. V., McElroy, S. L., Harnett-Sheehan, K., Keck, P. E., Janavs, J., Rogers, J., et al. (2009, June). Randomized, placebo-controlled trial of risperidone for acute treatment of bipolar anxiety. *Journal of Affective Disorders*, 115(3), 376-385.
- Shelledy, D. C., Legrand, T. S., Gardner, D. D., & Peters, J. I. (2009, March). A randomized, controlled study to evaluate the role of an in-home asthma disease management program provided by respiratory therapists in improving outcomes and reducing the cost of care. *Journal of Asthma*, 46(2), 194-201.
- Shepperd, S., Doll, H., Angus, R. M., Clarke, M. J., Iliffe, S., Kalra, L., et al. (2009, January 20). Avoiding hospital admission through provision of hospital care at home: a systematic review and meta-analysis of individual patient data. *Canadian Medical Association Journal*, 180(2), 175-182.
- Shepperd, S., Doll, H., Broad, J., Gladman, J., Iliffe, S., Langhorne, P., et al. (2009). Early discharge hospital at home. *Cochrane Database of Systematic Reviews*(1).
- Simon, G. E., Von Korff, M., & Lin, E. (2005, February). Clinical and functional outcomes of depression treatment in patients with and without chronic medical illness. *Psychological Medicine*, 35(2), 271-279.
- Sirtori, V., Corbetta, D., Moja, L., & Gatti, R. (2009). Constraint-induced movement therapy for upper extremities in stroke patients. *Cochrane Database of Systematic Reviews*(4).
- Skumlien, S., Skogedal, E. A., Skrede Ryg, M., & Bjørtuft, Ø. (2008, March). Endurance or resistance training in primary care after in-patient rehabilitation for COPD? *Respiratory Medicine*, 102(3), 422-429.
- Slade, M., Taber, D., Clarke, M. M., Johnson, C., Kapoor, D., Leikin, J. B., et al. (2007, November-December). Best practices for the treatment of patients with mental and substance use illnesses in the emergency department. *Disease-a-Month*, 53(11-12), 536-580.
- Slapø, G. D., Lossius, M. I., & Gjerstad, L. (2006, December). Poststroke epilepsy: occurrence, predictors and treatment. *Expert Review of Neurotherapeutics*, 6(12), 1801-1809.
- Smelson, D. A., Dixon, L., Craig, T., & Remolina, S. (2008). Pharmacological Treatment of Schizophrenia and Co-Occurring Substance Use Disorders. *CNS Drugs*, 22(11), 903-916.
- Smeulders, E. S., van Haagstregt, J. C., Ambergen, T., Janssen-Boyne, J. J., van Eijk, J. T., & Kempen, G. I. (2009, June). The impact of a self-management group programme on health behaviour and healthcare utilization among congestive heart failure patients. *European Journal of Heart Failure*, 11(6), 609-616.
- Smith, S.A., Shah, N.D., Bryant, S.C., Christianson, T.J.H., Bjornsen, S.S., Giesler, P.D., et al. (2008, July). Chronic Care Model and Shared Care in Diabetes: Randomized Trial of an Electronic Decision Support System. *Mayo Clinic Proceedings*, 83(7), 747-757.
- Sobell, L. C., Sobell, M. B., & Agrawal, S. (2009, December). Randomized controlled trial of a cognitive-behavioral motivational intervention in a group versus individual format for substance use disorders. *Psychology of Addictive Behaviors*, 23(4), 672-683.

Song, M.K., Ward, S. E., Happ, M. B., Piraino, B., Donovan, H. S., Shields, A.M., et al. (2009, June). Randomized controlled trial of SPIRIT: An effective approach to preparing African-American dialysis patients and families for end of life. *Research in Nursing & Health*, 32(3), 260-273.

Soran, O. Z., Piña, I. L., Lamas, G. A., Kelsey, S. F., Selzer, F., Pilotte, J., et al. (2008, November). A Randomized Clinical Trial of the Clinical Effects of Enhanced Heart Failure Monitoring Using a Computer-Based Telephonic Monitoring System in Older Minorities and Women. *Journal of Cardiac Failure*, 14(9), 711-717.

Spinewine, A., Swine, C., Dhillon, S., Lambert, P., Nachega, J. B., Wilmotte, L., et al. (2007, May). Effect of a collaborative approach on the quality of prescribing for geriatric inpatients: a randomized, controlled trial. *Journal of the American Geriatrics Society*, 55(5), 658-665.

Spruit, M. A., Pennings, H. J., Janssen, P. P., Does, J. D., Scroyen, S., Akkermans, M. A., et al. (2007, December). Extra-pulmonary features in COPD patients entering rehabilitation after stratification for MRC dyspnea grade. *Respiratory Medicine*, 101(12), 2454-2463.

Starkstein, S. E., Jorge, R., Petracca, G., & Robinson, R. G. (2007, January). The Construct of Generalized Anxiety Disorder in Alzheimer Disease. *The American Journal of Geriatric Psychiatry*, 15(1), 42-49.

Stephen J Bartels, E. H., Bartels, S. J., Coakley, E. H., Zubritsky, C., & Ware, J. H. (2004, August). Improving Access to Geriatric Mental Health Services: A Randomized Trial Comparing Treatment Engagement With Integrated Versus Enhanced Referral Care for Depression, Anxiety, and At-Risk Alcohol Use. *The American Journal of Psychiatry*, 161(8), 1455-1463.

Steuten, L., Vrijhoef, B., Van Merode, F., Wesseling, G.J., & Spreeuwenberg, C. (2006, December). Evaluation of a regional disease management programme for patients with asthma or chronic obstructive pulmonary disease. *International Journal for Quality in Health Care*, 18(6), 429-436.

Stewart, M., Barnfather, A., Neufield, A., Warren, S., Letourneau, N., & Liu, L. (2006, Summer). Accessible support for family caregivers of seniors with chronic conditions: from isolation to inclusion. *Canadian Journal on Aging*, 25(2), 179-192.

Stewart, S. H., Latham, P. K., Miller, P. M., Randall, P., & Anton, R. F. (2008, October). Blood pressure reduction during treatment for alcohol dependence: results from the Combining Medications and Behavioral Interventions for Alcoholism (COMBINE) study. *Addiction*, 103(10), 1622-1628.

Stiefel, F., Zdrojewski, C., Bel Hadj, F., Boffa, D., Dorogi, Y., So, A., et al. (2008). Effects of a multifaceted psychiatric intervention targeted for the complex medically ill: a randomized controlled trial. *Psychotherapy and Psychosomatics*, 77(4), 247-256.

Stimmel, G.L. (2001). Maximizing Treatment Outcome in Depression: Strategies to Overcome Social Stigma and Noncompliance. *Disease Management & Health Outcomes*, 9(4), 179-186.

Strippoli, G. F., Bonifati, C., Craig, M. E., Navaneethan, S. D., & Craig, J. C. (2006). Angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists for preventing the progression of diabetic kidney disease. *Cochrane Database of Systematic Reviews*(4).

Strippoli, G. F., Craig, M. E., Craig, J. C., Lv, J., & Perkovic, V. (2005, October 19). Antihypertensive agents for preventing diabetic kidney disease. *Cochrane Database of Systematic Reviews*(4).

Stuck, A. E., Egger, M., Hammer, A., Minder, C. E., & Beck, J. C. (2002, February 27). Home visits to prevent nursing home admission and functional decline in elderly people: systematic review and meta-regression analysis. *The Journal of the American Medical Association*, 287(8), 1022-1028.

Swartz, M. S., Wagner, H. R., Swanson, J. W., Stroup, S., McEvoy, J. P., Reimherr, F., et al. (2008, March). The effectiveness of antipsychotic medications in patients who use or avoid illicit substances: Results from the CATIE study. *Schizophrenia Research*, 100(1-3), 39-52.

T

Taylor, B., Irving, H. M., Baliunas, D., Roerecke, M., Patra, J., Mohapatra, S., et al. (2009, December). Alcohol and hypertension: gender differences in dose-response relationships determined through systematic review and meta-analysis. *Addiction*, 104(12), 1981-90.

Taylor, S. J., Bestall, J. C., Cotter, S., Falshaw, M., Hood, S. G., Parsons, S., et al. (2005). Clinical service organisation for heart failure. *Cochrane Database of Systematic Reviews*(2).

Thorpe, L., & Groulx, B. (2001, May). Depressive Syndromes in Dementia. *Canadian Journal of Neurological Sciences*, 28(Supplement 1), 83-95.

Tinetti, M. E., Baker, D., Gallo, W. T., Nanda, A., Charpentier, P., & O'Leary, J. (2002, April 24). Evaluation of restorative care vs usual care for older adults receiving an acute episode of home care. *The Journal of the American Medical Association*, 287(16), 2098-2105.

Tobe, S. W., Pylpchuk, G., Wentworth, J., & Kiss, A. (2006, April 25). Effect of nurse-directed hypertension treatment among First Nations people with existing hypertension and diabetes mellitus: the Diabetes Risk Evaluation and Microalbuminuria (DREAM 3) randomized controlled trial. *Canadian Medical Association. Journal*, 174(9), 1267-1271.

Tohen, M., Calabrese, J., Vieta, E., Bowden, C., Gonzalez-Pinto, A., Lin, D., et al. (2007, December). Effect of comorbid anxiety on treatment response in bipolar depression. *Journal of Affective Disorders*, 104(1-3), 137-146.

Troosters, T., Gosselink, R., & Decramer, M. (2004, May-June). Chronic obstructive pulmonary disease and chronic heart failure: two muscle diseases? *Journal of Cardiopulmonary Rehabilitation*, 24(3), 137-145.

U

Unützer J., Katon, W., Callahan, C.M., Williams, J.W., Hunkeler, E., Harpole, L., Hoffing, M., et al. (2002). Collaborative Care Management of Late-Life Depression in the Primary Care Setting - A Randomized Controlled Trial. *The Journal of the American Medical Association*, 288, 2836-2845.

V

van der Meer, R. M., Wagena, E., Ostelo, R., Jacobs, A. J., & van Schayck, O. P. (2003). Smoking cessation for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*(2).

van Eijk, J. T., Diederiks, J. P., Kempen, G. I., Honig, A., van der Meer, K., & Brenninkmeijer, W. J. (2004, July). Development and feasibility of a nurse administered strategy on depression in community-dwelling patients with a chronic physical disease. *Patient Education & Counseling*, 54(1), 87-94.

van Straten, A., Tiemans, B., Hakkaart, L., Nolen, W. A., & Donker, M. C. (2006, June). Stepped care vs. matched care for mood and anxiety disorders: a randomized trial in routine practice. *Acta Psychiatrica Scandinavica*, 113(6), 468-476.

Vanelle, J. M., & Douki, S. (2006, December). A double-blind randomised comparative trial of amisulpride versus olanzapine for 2 months in the treatment of subjects with schizophrenia and comorbid depression. *European Psychiatry*, 21(8), 523-530.

Veazey, C., Cook, K. F., Stanley, M., Lai, E. C., & Kunik, M. E. (2009, September). Telephone-administered cognitive behavioral therapy: a case study of anxiety and depression in Parkinson's disease. *Journal of Clinical Psychology in Medical Settings*, 16(3), 243-253.

Vernooij-Dassen, M., & Olde Rikkert, M. G. (2004, August). Personal disease management in dementia care. *International Journal of Geriatric Psychiatry*, 19(8), 715-717.

Vickrey, B. G., Mittman, B. S., Connor, K. I., Pearson, M. L., Della Penna, R. D., Ganiats, T. G., et al. (2006, November 21). The effect of a disease management intervention on quality and outcomes of dementia care: a randomized, controlled trial. *Annals of Internal Medicine*, 145(10), 713-726.

Von Muenster S.J., Carter B.L., Weber C.A., Ernst M.E., Milchak J.L., Steffensmeier J.J., Xu Y. (2008, January). Description of pharmacist interventions during physician-pharmacist co-management of hypertension. *Pharmacy World & Science*, 30(1),128-135.

Vrijhoef, H. J., Van Den Bergh, J. H., Diederiks, J. P., Weemhoff, I., & Spreeuwenberg, C. (2007, June). Transfer of care for outpatients with stable chronic obstructive pulmonary disease from respiratory care physician to respiratory nurse--a randomized controlled study. *Chronic Illness*, 3(2), 130-144.

W

- Wakefield, B. J., Ward, M. M., Holman, J. E., Ray, A., Scherubel, M., Burns, T. L., et al. (2008, October). Evaluation of home telehealth following hospitalization for heart failure: a randomized trial. *Telemedicine Journal and e-health*, 14(8), 753-761.
- Waldemar, G., Dubois, B., Emre, M., Georges, J., McKeith, I. G., Rossor, M., et al. (2007, January). Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *European Journal of Neurology*, 14(1), e1-26.
- Walters, J. A., Turnock, A. C., Walters, E. H., & Wood-Baker, R. (2010). Action plans with limited patient education only for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*(5).
- Wanner, C., Krane, V., März, W., Olschewski, M., Mann, J. F., Ruf, G., et al. (2005, July 21). Atorvastatin in patients with type 2 diabetes mellitus undergoing hemodialysis. *New England Journal of Medicine*, 353(3), 238-248.
- Watkins, K. E., Paddock, S. M., Zhang, L., & Wells, K. B. (2006, January). Improving care for depression in patients with comorbid substance misuse. *American Journal of Psychiatry*, 163(1), 125-132.
- Weir, M. R., Yeh, F., Silverman, A., Devereux, R. B., Galloway, J. M., Henderson, J. A., et al. (2009, October). Safety and feasibility of achieving lower systolic blood pressure goals in persons with type 2 diabetes: the SANDS trial. *Journal of Clinical Hypertension*, 11(10), 540-548.
- Weisner, C., Mertens, J., Parthasarathy, S., Moore, C., & Lu, Y. (2001). Randomized trial of integrated primary medical care with addiction treatment. Positive health outcomes and reduced costs. *Journal of the American Medical Association*, 286, 1715-1723.
- Weiss, R. D., Griffin, M. L., Jaffee, W. B., Bender, R. E., Graff, F. S., Gallop, R. J., et al. (2009, October 1). A "community-friendly" version of integrated group therapy for patients with bipolar disorder and substance dependence: a randomized controlled trial. *Drug and Alcohol Dependence*, 104(3), 212-219.
- West, D. S., DiLillo, V., Bursac, Z., Gore, S. A., & Greene, P. G. (2007, May). Motivational interviewing improves weight loss in women with type 2 diabetes. *Diabetes Care*, 30(5), 1081-1087.
- Whellan, D. J., Hasselblad, V., Peterson, E., O'Connor, C. M., & Schulman, K. A. (2005, April). Metaanalysis and review of heart failure disease management randomized controlled clinical trials. *American Heart Journal*, 149(4), 722-729.
- Whitlock, E. P., Polen, M. R., Green, C. A., Orleans, T., & Klein, J. (2004, April 6). Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Annals of Internal Medicine*, 140(7), 557-568.

- Whitten, P., & Mickus, M. (2007). Home telecare for COPD/CHF patients: outcomes and perceptions. *Journal of Telemedicine and Telecare*, 13(2), 69-73.
- Whooley, M.A., de Jonge, P., Vittinghoff, E., Otte, C., Moos, R., Carney, R.M., Ali, S., Dowray, S., et al. (2008). Depressive Symptoms, Health Behaviors, and Risk of Cardiovascular Events in Patients with Coronary Heart Disease. *The Journal of the American Medical Association*, 300(20), 2379-2388.
- Williams, L. S., Kroenke, K., Bakas, T., Plue, L. D., Brizendine, E., Tu, W., et al. (2007, March). Care management of poststroke depression: a randomized, controlled trial. *Stroke*, 38(3), 998-1003.
- Williams, M. A., Fleg, J. L., Ades, P. A., Chaitman, B. R., Miller, N. H., Mohiuddin, S. M., et al. (2002, April 9). Secondary prevention of coronary heart disease in the elderly (with emphasis on patients > or =75 years of age): an American Heart Association scientific statement from the Council on Clinical Cardiology Subcommittee on Exercise, Cardiac Rehabilitation. *Circulation*, 105(14), 1735-1743.
- Williams P.T., Franklin B. (2007, November). Vigorous exercise and diabetic, hypertensive, and hypercholesterolemia medication use. *Medicine and Science in Sports and Exercise*, 39(11), 1933-1941.
- Wilson, D., Ipser, J. C., & Stein, D. J. (2008, October 8). Pharmacotherapy for anxiety disorders and comorbid alcohol dependency (Protocol). *Cochrane Database of Systematic Reviews*(4).
- Winter, L., Corcoran, M., Schinfeld, S., Gitlin, L. N., Dennis, M. P., & Hauck, W. W. (2006, May). A Randomized Trial of a Multicomponent Home Intervention to Reduce Functional Difficulties in Older Adults. *Journal of the American Geriatrics Society*, 54(5), 809-816.
- Winter, L., Schulz, R., Gitlin, L. N., Hauck, W. W., & Dennis, M. P. (2006, June). Effect of an In-Home Occupational and Physical Therapy Intervention on Reducing Mortality in Functionally Vulnerable Older People: Preliminary Findings. *Journal of the American Geriatrics Society*, 54(6), 950-955.
- Woo, J., Hui, E., Hui, D., Lum, C. M., Or, K. H., & Kwok, T. (2009, February). A pilot study to examine the feasibility and acceptability of a community model for exercise prescription for patients with chronic disease. *Hong Kong Medical Journal*, 15(Supplement 2), 12-16.
- Wright, J. M., & Musini, V. M. (2009). First-line drugs for hypertension. *Cochrane Database of Systematic Reviews*(3).
- Wu, J. Y., Leung, W. Y., Chang, S., Lee, B., Zee, B., Tong, P. C., et al. (2006, September 9). Effectiveness of telephone counselling by a pharmacist in reducing mortality in patients receiving polypharmacy: randomised controlled trial. *British Medical Journal*, 333(7567), 522.

Y

Yang, Y., Bellack, A. S., Bennett, M. E., Gearon, J. S., & Brown, C. H. (2006, April). A Randomized Clinical Trial of a New Behavioral Treatment for Drug Abuse in People With Severe and Persistent Mental Illness. *Archives of General Psychiatry*, 63(4), 426-432.

Yatham, L. N., Kennedy, S. H., O'Donovan, C., Parikh, S., MacQueen, G., McIntyre, R., et al. (2005). Canadian Network for Mood and Anxiety Treatments (CANMAT) guidelines for the management of patients with bipolar disorder: consensus and controversies. *Bipolar Disorders*, 7(Supplement 3), 5-69.

Yorke, J., Fleming, S. L., & Shuldham, C. M. (2006). Psychological interventions for adults with asthma. *Cochrane Database of Systematic Reviews*(1).

Young L.H., Wackers F.J., Chyun D.A., Davey J.A., Barrett E.J., Taillefer R., Heller G.V., Iskandrian A.E., Wittlin S.D., Filipchuk N., Ratner R.E., & Inzucchi S.E. (2009, April). Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: a randomized controlled trial. *The Journal of the American Medical Society*, 301(15), 1547-1555.

Z

Zaccara, G. (2009, July). Neurological comorbidity and epilepsy: implications for treatment. *Acta Neurologica Scandinavica*, 120(1), 1-15.

Zanchetti, A., & Ruilope, L. M. (2002, November). Antihypertensive treatment in patients with type-2 diabetes mellitus: what guidance from recent controlled randomized trials? *Journal of Hypertension*, 20(11), 2099-2110.

Zhang, N. J., Wan, T. T., Rossiter, L. F., Murawski, M. M., & Patel, U. B. (2008, May). Evaluation of chronic disease management on outcomes and cost of care for Medicaid beneficiaries. *Health Policy*, 86(2-3), 345-354.

Zhu K., Devine A., Dick I.M., Prince R.L. (2007, August 15). Association of back pain frequency with mortality, coronary heart events, mobility, and quality of life in elderly women. *Spine*, 32(18), 2012-2018.

Ziedonis, D. M., Smelson, D., Rosenthal, R. N., Batki, S. L., Green, A. I., Henry, R. J., et al. (2005, September). Improving the care of individuals with schizophrenia and substance use disorders: consensus recommendations. *Journal of Psychiatric Practice*, 11(5), 315-339.

Zisook, S., Montross, L., Kasckow, J., Mohamed, S., Palmer, B. W., Patterson, T. L., et al. (2007, December). Subsyndromal depressive symptoms in middle-aged and older persons with schizophrenia. *American Journal of Geriatric Psychiatry*, 15(12), 1005-1014.